

A nonsurgical, injectable option for adults with moderate to severe submental fat (double chin):

# Where do you draw the line?

A chin profile line that slopes down could affect your appearance.

DAVID, AGE: 53



BEFORE



AFTER 4 TREATMENTS

Unretouched photos of paid model. Individual results may vary.

Not all treatments are shown; 59% of adults received 6 KYBELLA<sup>®</sup> treatments in clinical studies.

**Learn more about improving your chin profile nonsurgically—permanently destroying the fat under your chin.**

## What is KYBELLA<sup>®</sup>?

KYBELLA<sup>®</sup> is a prescription medicine used in adults to improve the appearance and profile of moderate to severe fat below the chin (submental fat), also called “double chin.”

It is not known if KYBELLA<sup>®</sup> is safe and effective for the treatment of fat outside of the submental area or in children under 18 years of age.

## Important Safety Information

### Who should not receive KYBELLA<sup>®</sup>?

Do not receive KYBELLA<sup>®</sup> if you have an infection in the treatment area.

**Please see additional Important Safety Information inside and accompanying full Prescribing Information.**

# Men like you draw a different line with KYBELLA<sup>®</sup>

**KYBELLA<sup>®</sup> is the first-of-its-kind, nonsurgical option for the moderate to severe fullness under your chin**

- Injectable
- Permanently destroys fat cells in the treated area
- Improves the chin profile

JASON, AGE: 36



BEFORE

AFTER 6 TREATMENTS

Unretouched photos of paid model. Individual results may vary.

Not all treatments are shown; 59% of adults received 6 KYBELLA<sup>®</sup> treatments in clinical studies.

## **Important Safety Information (continued)**

**Before receiving KYBELLA<sup>®</sup>, tell your healthcare provider about all of your medical conditions, including if you:** Have had or plan to have surgery on your face, neck, or chin; have had cosmetic treatments on your face, neck, or chin; have had or have medical conditions in or near the neck area; have had or have trouble swallowing; have bleeding problems; are pregnant or plan to become pregnant (it is not known if KYBELLA<sup>®</sup> will harm your unborn baby); are breastfeeding or plan to breastfeed (it is not known if KYBELLA<sup>®</sup> passes into your breast milk).

## **A series of treatments individually tailored\* to your unique chin profile**

You and your provider will determine the number of treatments you may need, based on:

- How much fullness is beneath your chin
- Your aesthetic goals

\*Multiple injections under the chin per treatment; up to 6 treatments at least 1 month apart.

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take a medicine that prevents the clotting of your blood (antiplatelet or anticoagulant medicine).

**Please see additional Important Safety Information throughout this brochure and accompanying full Prescribing Information.**

## Is your chin profile line sloping down?



Extra fat below the chin, also called double chin, could be affecting your appearance.



Look at a photo of yourself from the side.




How much does your profile line slope?

### Important Safety Information (continued) What are the possible side effects of KYBELLA®? KYBELLA® can cause serious side effects, including

- Nerve injury in the jaw (which can cause an uneven smile or facial muscle weakness)
- Trouble swallowing

## Other men are bothered too



In an online survey\*

The **double chin** was reported as the **#2** facial area of concern for men

The double chin received **first priority** for aesthetic treatment by **7 in 10** men

\*Online survey of injectable-naïve, aesthetically oriented men aged 30 to 65 (N = 600).

## Diet and exercise alone may not help—submental fullness can be caused by more than weight

Potential causes of submental fullness



Weight Gain



Genetics



Aging

## Ask for a chin profile assessment today. See if KYBELLA® is right for you.

- Injection site problems including: a collection of blood under the skin (hematoma) or bruising, damage to an artery or vein if KYBELLA® is inadvertently injected into it, hair loss, open sores (ulcers), damage and tissue cell-death (necrosis) around the injection site. Call your healthcare provider if you: begin to develop weakness in the muscles of your face, or your smile becomes uneven; have difficulty swallowing, or if any of the symptoms that you already have get worse; develop open sores or drainage from the treatment area

Please see additional Important Safety Information throughout this brochure and accompanying full Prescribing Information.

## How does KYBELLA® work?

### Once the fat cells in the treatment area are gone, they're gone!

- Deoxycholic acid is a naturally occurring molecule that helps break down fat in the body
- The active ingredient in KYBELLA® is synthetic deoxycholic acid
- Fat cells are broken down each time KYBELLA® is injected into the treatment area beneath the chin
- Over a period of time, those cells are cleared away through the natural processes of your body
- You are left with **a noticeable reduction in fullness under the chin**

EVALUATED IN  
APPROXIMATELY  
**20** CLINICAL  
STUDIES  
AROUND THE WORLD

APPROXIMATELY  
**1600** PEOPLE  
WORLDWIDE  
WERE TREATED  
WITH KYBELLA®

#### Important Safety Information (continued)

#### What are the possible side effects of KYBELLA®? (continued)

The most common side effects of KYBELLA® include swelling, pain, numbness, redness, and areas of hardness in the treatment area.

These are not all of the possible side effects of KYBELLA®. Call your doctor for medical advice about side effects.

Please see additional Important Safety Information throughout this brochure and accompanying full Prescribing Information.

## Progressive results you can see

Seeing results is different for everyone.

NIKKI, AGE: 35



BEFORE




AFTER 2 TREATMENTS

JASON, AGE: 36




BEFORE



AFTER 2 TREATMENTS



AFTER 4 TREATMENTS



AFTER 6 TREATMENTS

Unretouched photos of paid models. Individual results may vary.

Not all treatments are shown; 59% of adults received 6 KYBELLA® treatments in clinical studies.


**It is important to talk to your provider  
about what you can expect**

## Noticeable results for an improved chin profile

ADRA, AGE: 35



BEFORE




AFTER 3 TREATMENTS

Unretouched photos of paid models. Individual results may vary.  
Not all treatments are shown; 59% of adults received 6 KYBELLA® treatments in clinical studies.

DAVID, AGE: 53



BEFORE



AFTER 4 TREATMENTS

ANDY, AGE: 46




BEFORE




AFTER 4 TREATMENTS

JASON, AGE: 36



BEFORE



AFTER 6 TREATMENTS

### Important Safety Information (continued)

#### Who should not receive KYBELLA®?

Do not receive KYBELLA® if you have an infection in the treatment area.

#### Before receiving KYBELLA®, tell your healthcare provider about all of your medical conditions, including if you:

Have had or plan to have surgery on your face, neck, or chin; have had cosmetic treatments on your face, neck, or chin; have had or have medical conditions in or near the neck area; have had or have

trouble swallowing; have bleeding problems; are pregnant or plan to become pregnant (it is not known if KYBELLA® will harm your unborn baby); are breastfeeding or plan to breastfeed (it is not known if KYBELLA® passes into your breast milk).

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take a medicine that prevents the clotting of your blood (antiplatelet or anticoagulant medicine).

### What are the possible side effects of KYBELLA®?

#### KYBELLA® can cause serious side effects, including

- Nerve injury in the jaw (which can cause an uneven smile or facial muscle weakness)
- Trouble swallowing
- Injection site problems including: a collection of blood under the skin (hematoma) or bruising, damage to an artery or vein if KYBELLA® is inadvertently injected into it, hair loss, open sores (ulcers), damage and tissue cell-death (necrosis) around the injection site. Call your healthcare provider if you: begin

to develop weakness in the muscles of your face, or your smile becomes uneven; have difficulty swallowing, or if any of the symptoms that you already have get worse; develop open sores or drainage from the treatment area

**Please see additional Important Safety Information throughout this brochure and accompanying full Prescribing Information.**

## What to expect



### **Treat. Repeat. Complete.**

Your provider will recommend the number of treatment sessions you may need to achieve your aesthetic goals.



### **Each treatment is at least 1 month apart**

Schedule your next appointment before you leave the office.



### **Office staff will take your photo before and after each treatment**

Track the change in your chin profile line throughout your treatment series.

THE INJECTION PROCESS MAY TAKE  
**15 TO 20 MINUTES**



Your provider may use cold packs or local anesthesia to help make you more comfortable

## Possible side effects

Sometimes serious side effects can occur, including:

- Nerve injury in the jaw that can temporarily (median 44 days) cause an uneven smile or facial muscle weakness (experienced by 4 out of 100 people in clinical studies)
- Trouble swallowing (experienced by 2 out of 100 people in clinical studies)
- Possible injection-site problems include a collection of blood under the skin (hematoma) or bruising, damage to an artery or vein if KYBELLA<sup>®</sup> is inadvertently injected into it, hair loss, open sores (ulcers), damage and tissue cell-death (necrosis) around the injection site. Call your healthcare provider if you: begin to develop weakness in the muscles of your face, or your smile becomes uneven; have difficulty swallowing, or if any of the symptoms that you already have get worse; develop open sores or drainage from the treatment area

The most common side effects in the treatment area included:

- Swelling
- Numbness
- Redness
- Pain
- Areas of hardness

### **Swelling is an expected reaction to treatment**

It generally became less severe and happened less often with subsequent treatment sessions in studies.

**TIP:** When scheduling treatment, consider planned social or work commitments, as side effects like swelling are common.

### **Example of swelling after first treatment**



BEFORE

FIRST DAY AFTER TREATMENT 1

BEFORE TREATMENT 2

Individual patient experience. Swelling reactions may vary. 87% of adults receiving KYBELLA<sup>®</sup> treatment in clinical trials experienced swelling as a side effect.

Ask your provider about *treatment package pricing* to help you save, since multiple treatments are likely.

Watch real stories about the KYBELLA<sup>®</sup> experience.  
**Check out MyKybella.com**

# The first of its kind

- ✓ An injectable, nonsurgical option
- ✓ Permanently destroys fat cells under the chin
- ✓ Individually tailored\* to improve your chin profile

\*Multiple injections under the chin per treatment; up to 6 treatments at least 1 month apart.

JASON, AGE: 36



BEFORE



AFTER 6 TREATMENTS

Unretouched photos of paid model. Individual results may vary.

Not all treatments are shown; 59% of adults received 6 KYBELLA<sup>®</sup> treatments in clinical studies.

**Find out if KYBELLA<sup>®</sup> is right for you—ask for a chin profile assessment today.**



Earn savings and rewards on KYBELLA<sup>®</sup> and other leading aesthetic treatments from Allergan. Join today at [BrilliantDistinctionsProgram.com](http://BrilliantDistinctionsProgram.com).

## Important Safety Information (continued)

### What are the possible side effects of KYBELLA<sup>®</sup>? (continued)

The most common side effects of KYBELLA<sup>®</sup> include swelling, pain, numbness, redness, and areas of hardness in the treatment area.

These are not all of the possible side effects of KYBELLA<sup>®</sup>. Call your doctor for medical advice about side effects.

**Please see additional Important Safety Information throughout this brochure and full Prescribing Information inside the pocket of this brochure.**